

SANDWICH Checklist



Has your patient been set a COMFORT target?



Have you assessed your patient's COMFORT score regularly?



Have you completed the Bedside Screen this shift?



Is your patient ready for a SBT?



Is your patient ready to extubate?




Have you discussed all above on the MDT ward round?



Setting COMFORT targets

Origina



FLACCS < 4

Issue is SEDATION

Consider increasing/ adapting Sedation

FLACCS > 4

Issue is PAIN

Consider increasing/ adapting analgesia

COMFORT score too high. Complete FLACCS pain score to differentiate source of high score. Modify sedation and/or analgesia as recommended.



FLACCS < 4

Patient is comfortable

Optimal level of sedation for patients to breath up, cough spontaneously, respond to environment but not distressed / agitated / pulling lines or tubes etc



FLACCS < 4

Consider Weaning Sedation

Moderately suppressed reflexes- Slightly more sedate, less response to environment/stimuli but continues to trigger vent & cough to stimulation eg. Bronchiolitics



Patient is **OVER SEDATED**Wean Sedation

Very heavily sedated- not likely to trigger vent or cough spontaneously. Not responsive to environment or stimuli eg. Traumatic brain injury, ECHMO

Spontaneous mode of ventilation

PEEP of **5cm** H20 & Pressure Support of **5cm** H20 above PEEP